Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Ocular Oncology and Oculoplasty

Journal homepage: <https://ijooo.org/>

Case Report

Unusual presentation of oculosporeidiosis: A case report

Narayan Chandra Bhowmik^{1,*}, Syeed Mehbub Ul Kadir², Riffat Rashid³,
A B Shamsudduha¹, Ashraf Sayeed¹

¹Dept. of Ophthalmology, BIRDEM General Hospital, Dhaka, Bangladesh

²Dept. of Oculoplasty, Sheikh Fazilatunnessa Mujib Eye Hospital and Training Institute, Gopalganj, Dhaka, Bangladesh

³Dept. of Oculoplasty, Ispahani Islamia Eye Institute & Hospital, Dhaka, Bangladesh

ARTICLE INFO

Article history:

Received 05-10-2021

Accepted 15-10-2021

Available online 25-10-2021

Keywords:

Lacrimal sac

Oculosporeidiosis

Excision

Dapsone

ABSTRACT

Rhinosporidiosis is a chronic localized granulomatous disease caused by *Rhinosporidium seeberi*, an aquatic protistan parasite belonging to a clade, Mesomycetozoa. Infestation of *Rhinosporidiosis* on the eye and adnexa is termed oculosporeidiosis. In such cases, conjunctival mucosa is mostly involved. Our case was a 25 years old male, presented with swelling of the lacrimal sac and deeper periorbital tissue and was presented as a case of orbital mass. Complete surgical excision of the lesion was done and Histopathological examination confirms orbital rhinosporidiosis. Postoperatively dapsone therapy was given for 6 months and the patient responded very well with no recurrence during the 6 months follow-up period.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](#), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Rhinosporidiosis is a parasitic disease and presented as a chronic localized granulomatous disease caused by *Rhinosporidium seeberi*.¹ Rhinosporidiosis is a common disease and endemic in India as well as Sri Lanka but quite is sporadic in other parts of the world.^{2,3} Most common site of involvement is nasal mucosa, followed by ocular involvement. Other natural orifices such as the penis, urethra, anus, vagina, skin⁴ may also be involved; few cases of disseminated rhinosporidiosis have also been reported in the literature.^{5,6} Rhinosporidiosis of the eye and adnexa is termed as oculosporeidiosis.⁷ Clinically, oculosporeidiosis appears as a friable polypoid mass, causing a foreign body sensation, irritation and watering from the eye, and no effects on vision according to fewer case reports.⁸ Oculosporeidiosis commonly occurs in young and younger adults (15–40 years) with a predominance in males.⁹

Here, we present an interesting case of oculosporeidiosis. How the patient was diagnosed and treated is also provided, along with the recent literature review.

2. Case Report

A 25-year-old male patient was presented with a swelling over the right lower periorbital area for the last 3 years. On examination, the swelling was pinkish in color with an oblique linear scar over it due to previous surgical intervention [Figure 1]. On palpation, the swelling was soft, nontender, and fixed with the deeper tissue.

Two months ago, he was treated as a case of orbital abscess of his right eye for which FNAC was done, then incision and drainage were made by an ophthalmologist in his local area; it took around 2 weeks for the wound to heal. Although there was an initial reduction of the size of the swelling for a month, the swelling started to grow again for the last one month.

* Corresponding author.

E-mail address: dr.narayan4@gmail.com (N. C. Bhowmik).

Fig. 1: **a:** The patient presented right orbital swelling with a scar on it after two months of primary treatment. **b:** The patient was initially introduced on the right lower orbital lesion diagnosed as orbital abscess and drained the abscess locally two months ago.

He was treated at the tertiary-based eye hospital, Dhaka. CT Scan of orbit was done and nasal swabs were collected. Contrast-enhanced computed tomography scan of the orbit showed soft tissue lesion (3.2 x2.3x1.6 mm) with intralesional loculated fluid collection in the inferomedial aspect of the right orbit. Mild heterogeneous enhancement of the lesion compresses the right eyeball [Figure 2 a,b]. The impression of the radiologist was an inflammatory lesion of the lacrimal sac with pyoceles and bilateral maxillary sinusitis.

Fig. 2: **a,b:** CT scan of orbit showing soft tissue mass in the lacrimal sac fossa with a bony breach and extension into inferomedial aspect of orbit; **c:** Repeat CT scan of orbit showing enhancing soft tissue mass in the floor of the right orbit.

Repeat incision and drainage were conducted by an ophthalmologist at that hospital. After external healing of the wound, the swelling started to grow again for the last one month. After that, he came to us (Dept. of ophthalmology, BIRDEM General Hospital) for treatment.

On examination, his visual acuity was found to be normal; conjunctiva was normal; syringing through both puncta of his right upper and lower eyelid proved the patency of the lacrimal drainage system. Repeat CT scan of orbit was done which showed enhancing two soft tissue masses in the floor of the right orbit measuring about (1.7x1.4) and (3.2x1.2) cm with thickening of perilesional fat. [Figure 2c]

His routine blood investigation along with erythrocyte sedimentation rate was normal. The patient was tested

negative for the COVID-19 test. His X-ray chest was done for a routine pre-anesthesia check-up, which was found to be normal.

Hence, with a preoperative diagnosis of orbital swelling due to lacrimal sac pyoceles, we decided to explore the area surgically. The operation was performed under general anesthesia. An incision was placed over the old scar extending medially up to the medial canthal ligament. On exploration, we found an ill-defined soft tissue mass adherent with the surrounding soft tissues; meticulous dissection was done and the swelling was found to be extending deep into the orbit. As the lacrimal sac was closely adherent with the mass, it was removed. [Figure 3a,b]. Extra-ocular muscles were not involved and intraoperative tests were done to check their functions. An Electro-cauterization of the wall of the cavity was done to achieve hemostasis; the wound was then closed.

Fig. 3: Intraoperative picture with specimen.

Fig. 4: Histopathology showing the typical sporangia containing numerous spores

Histopathological examination of the specimen was conducted. Sections from the orbital cyst and lacrimal sac showed tissue-lined partially by stratified squamous epithelium and partially by columnar epithelium. Sub-epithelium showed sporangia with endospores. Surrounding

tissues showed chronic inflammatory cell infiltrate [Figure 4]. Thus, a diagnosis of orbital rhinosporidiosis was made.

The patient had early recovery and was discharged on the 3rd postoperative day with oral antimicrobial therapy (dapsone 100 mg/day, for 6 days a week, for 6 months). Before discharge, a thorough ENT check-up was done with a nasal endoscope and fiberoptic laryngoscope to look for any such lesion in the upper respiratory passage, but no such lesion indicating infestation by *Rhinosporidium* in that area was found. There was no sign of recurrence in the last 6 months follow-up. [Figure 5]

Fig. 5: Postoperative (Patient on 6 months follow up).

3. Discussion

Oculosporidiosis is usually found among 10-15% of rhinosporidiosis cases. In the cases of oculosporidiosis, the commonest affected tissue is conjunctiva (77.6%), followed by lacrimal sac (26%) with or without conjunctiva involvement. Kuriakose ET described the clinical features of the lacrimal sac oculosporidiosis, as it may present with a painless, soft, swelling in the lower eyelid; which is fluctuating to the touch; there may be a history of bleeding from the nose, and haemolacria is seldom completely obstructed.⁸

In our case, though there was no bleeding from the nose, other features were present. But still, we missed the diagnosis because isolated oculosporidiosis of the lacrimal sac is very rare. So it becomes challenging to suspect a secluded lacrimal sac is swelling to be due to infestation by *Rhinosporidium*.¹⁰ They described a patient with recurrent painful swelling of the right lower eyelid, who was diagnosed elsewhere as a case of chronic dacryocystitis. They diagnosed it as a case of idiopathic orbital inflammatory disease.¹¹ Final diagnosis was made

only after a tissue biopsy from the lacrimal sac area.

A CT scan of the orbit may reveal an enlarged lacrimal sac, along with polypus mass inside the sinuses, thus helping to diagnose the lesions of rhinosporidiosis. However, in this case, the CT scan of the orbit showed enhanced heterogenous lesion involving the right lacrimal sac that showed a lacrimal abscess, paranasal sinuses were revealed as normal study. Thus, he was diagnosed with a chance of lacrimal sac pyocele. However, in suspected cases, the histopathological examination can confirm this disease, which shows characteristic features as numerous sporocysts in various stages of maturation.

Complete surgical excision is the mainstay of the Management of lacrimal oculosporidiosis;⁵ but there is a very high chance of recurrence, as reported by two large case series.⁷ The recurrence may occur due to incomplete removal of the infested sac.^{10–12}

It is often difficult to dissect the involved lacrimal sac becomes because of the risk of severe intraoperative bleeding.¹³ In our case, we experienced the same, and the bleeding was controlled by gel foam application. After excision of the lesion, copious irrigation with 5% povidone-iodine solution and 5 mg/ml amphotericin-B have also been tried with good results.¹²

Adjuvant antimicrobial dapsone therapy is used as an additional measure to reduce the possibility of recurrence as it has been found to halt the maturation of spores and promote fibrosis in the stroma.^{12,14} Although, some study suggests that dapsone therapy may not be very effective.¹⁵

In association with surgery, dapsone may be used to prevent the maturation of sporangia and promote fibrosis.¹⁶ Histopathological analysis revealed the confirmed diagnosis, with a demonstration of a fibrous stroma with several thick-walled sporangia at different stages of maturation and containing numerous endospores.^{17,18}

4. Conclusion

Isolated oculosporidiosis involving the lacrimal sac and presenting as a periorbital swelling is a rare clinical finding that can initially be misdiagnosed. In this case, it was wrongly diagnosed as a pyocele. The lesion, in such cases, must be surgically excised. Surgical excision followed by histopathological examination is vital for any suspicious lacrimal sac lesion. Postoperatively, dapsone therapy may be an additional measure to reduce the possibility of recurrence.

5. Conflict of Interest

The authors declare that there are no conflicts of interest in this paper.

6. Source of Funding

None.

References

- Herr RA, Ajello L, Taylor JW, Arseculeratne SN, Mendoza L. Phylogenetic analysis of Rhinosporidium seeberi's 18S small-subunit ribosomal DNA groups this pathogen among members of the protostistan Mesomycetozoa clade. *J Clin Microbiol*. 1999;37(9):2750–4. doi:10.1128/JCM.37.9.2750-2754.1999.
- Arseculeratne SN. Recent advances in rhinosporidiosis and rhinosporidium seeberi. *Ind J Med Microbiol*. 2002;20(3):119–31.
- Shrestha SP, Hennig A, Parija SC. Prevalence of rhinosporidiosis of the eye and its adnexa in Nepal. *Am J Trop Med Hyg*. 1998;59(2):2331–4. doi:10.4269/ajtmh.1998.59.231.
- Ahluwalia KB, Maheshwari N, Deka RC. Rhinosporidiosis: A study that resolves etiologic controversies. *Am J Rhinol*. 1997;11(6):479–83. doi:10.2500/105065897780914938.
- Amritanand R, Nithyananth M, Cherian VM, Venkatesh K, Shah A. Disseminated rhinosporidiosis destroying the talus: A case report. *J Orthop Surg*. 2008;16(1):99–101. doi:10.1177/230949900801600123.
- Sood N, Agarwal MC, Gujani HC. Ocular rhinosporidiosis: A case report from Delhi. *J Infect Dev Ctries*. 2012;6(11):825–7. doi:10.3855/jidc.2397.
- Kuriakose ET. Oculosporeidiosis: Rhinosporidiosis of the eye. *Br J Ophthalmol*. 1963;47(6):346–9. doi:10.1136/bjo.47.6.346..
- Nair AG, Ali MJ, Kaliki S, Naik MN. Rhinosporidiosis of the tarsal conjunctiva. *Indian J Ophthalmol*. 2015;63(5):462–3. doi:10.4103/0301-4738.159897.
- Nuruddin M, Mudhar HS, Osmani M, Roy SR. Lacrimal sac rhinosporidiosis: clinical profile and surgical management by modified dacryocystorhinostomy. *Orbit*. 2014;33(1):29–32.
- Shrestha SP, Hennig A, Parija SC. Prevalence of rhinosporidiosis of the eye and its adnexa in Nepal. *Am J Trop Med Hyg*. 1998;59(2):231–4. doi:10.4269/ajtmh.1998.59.231.
- Mukherjee B, Mohan A, Sumathi V, Biswas J. Infestation of the lacrimal sac by Rhinosporidium seeberi: A clinicopathological case report. *Indian J Ophthalmol*. 2013;61(10):588–90. doi:10.4103/0301-4738.121084.
- Ghorpade A, Gurumurthy J, Banerjee PK, Banerjee AK, Bhalla M, Ravindranath M, et al. Oculosporeidiosis presenting as an under-eye swelling. *Indian J Dermatol Venereol Leprol*. 2007;73(3):196–203.
- Chowdhury RK, Behera S, Bhuyan D, Das G. Oculosporeidiosis in a tertiary care hospital of western Orissa, India: A case series. *Indian J Ophthalmol*. 2007;55(4):299–301. doi:10.4103/0301-4738.33045..
- Sood A, Badhu B. Oculosporeidiosis in a tertiary care hospital of Western Orissa, India. *Indian J Ophthalmol*. 2008;56(2):165. doi:10.4103/0301-4738.39130.
- Job A, Venkateswaran S, Mathan M, Krishnaswami H, Raman R. Medical therapy of rhinosporidiosis with dapsone. *J Laryngol Otol*. 1993;107(9):809–12. doi:10.1017/s002221510012448x.
- Banjara H, Panda RK, Daharwal AV, Sudarshan V, Singh D, Gupta A, et al. Bronchial rhinosporidiosis: An unusual presentation. *Lung India*. 2012;29(2):173–5. doi:10.4103/0970-2113.95336..
- Shrestha SP, Hennig A, Parija SC. Prevalence of rhinosporidiosis of the eye and its adnexa in Nepal. *Am J Trop Med Hyg*. 1998;59(2):231–4.
- Prabhu SM, Irodi A, Kiangte HL, Rupa V, Naina P. Imaging features of Rhinosporidiosis on contrast CT. *Indian J Radiol Imaging*. 2013;23(3):212–18. doi:10.4103/0971-3026.120267.

Author biography

Narayan Chandra Bhowmik, Assistant Professor

Syed Mehbub Ul Kadir, Assistant Professor

Riffat Rashid, Associate Professor

A B Shamsudduha, Assistant Professor

Ashraf Sayeed, Professor and Head

Cite this article: Bhowmik NC, Kadir SMU, Rashid R, Shamsudduha AB, Sayeed A. Unusual presentation of oculosporeidiosis: A case report. *IP Int J Ocul Oncol Oculoplasty* 2021;7(3):299–302.